

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PROCESSES FOR PREPARATION OF 9,11-EPOXY
STEROIDS AND INTERMEDIATES USEFUL THEREINAbstract of the Disclosure

Multiple novel reaction schemes, novel process steps and novel intermediates are provided for the synthesis of epoxymexrenone and other compounds of Formula I




wherein:

-A-A- represents the group $-\text{CHR}^4-\text{CHR}^5-$ or $\text{CR}^4=\text{CR}^5-$

R^3 , R^4 and R^5 are independently selected from the group consisting of hydrogen, halo, hydroxy, lower alkyl, lower alkoxy, hydroxyalkyl, alkoxyalkyl, hydroxycarbonyl, cyano, aryloxy,

R^1 represents an alpha-oriented lower alkoxy carbonyl or hydroxyalkyl radical,

-B-B- represents the group $-\text{CHR}^6-\text{CHR}^7-$ or an alpha- or beta-oriented group:



where R^6 and R^7 are independently selected from

the group consisting of hydrogen, halo, lower alkoxy, acyl, hydroxyalkyl, alkoxyalkyl, hydroxycarbonyl, alkyl, alkoxy carbonyl, acyloxyalkyl, cyano, aryloxy, and

30

R^8 and R^9 are independently selected from the group consisting of hydrogen, halo, lower alkoxy, acyl, hydroxyalkyl, alkoxyalkyl, hydroxycarbonyl, alkyl, alkoxy carbonyl, acyloxyalkyl, cyano, aryloxy, or R^8 and R^9 together comprise a carbocyclic or heterocyclic ring structure, or R^8 or R^9 together with R^6 or R^7 comprise a carbocyclic or heterocyclic ring structure fused to the pentacyclic D ring.